

2. (a) Give an inductive definition of the set $A = \{3k + 5 : k \in \mathbb{N}\}$.

Basis:

Induction:

- (b) Give an inductive definition of the set C of all lists of odd length over $A = \{a, b\}$.

Basis:

Induction:

- (c) A set is defined inductively as follows:

Basis: $1 \in S$;

Induction: If $x \in S$, then $1 + \frac{1}{x} \in S$.

Write at least seven of the elements to give a flavor of S :

$$S = \{ \quad , \quad , \quad , \quad , \quad , \quad , \quad , \dots \}.$$

- (d) Give an inductive definition of the set

$$A = \{4, 7, 10, 12, \dots\} \times \{3, 9, 27, 81, \dots\}.$$

Basis:

Induction:

- (e) Write an inductive definition of $K = \{a^{2n} : n \in \mathbb{N}\} \cup \{b^{3n+2} : n \in \mathbb{N}\}$.

Basis:

Induction:

- (f) Consider the following set S of strings over alphabet $A = \{a, b\}$:

$$S = \{x \in \{a, b\}^* : x \text{ has the same number of } a\text{'s and } b\text{'s}\}.$$

The statement “ S is defined by the following inductive definition:

Basis: $\Lambda \in S$;

Induction: If $x \in S$, then $abx, bax, axb, bxa \in S$.”

is _____

Proof:

3. (a) Give a recursive definition of the function $\text{length} : \{a, b\}^* \rightarrow \mathbb{N}$, defined by

$$f(x) = \text{the length of } x, \quad \text{for all } x \in \{a, b\}^*.$$

Basis:

Recursion:

- (b) Give a recursive definition of the function $\text{dist} : A \times \text{Lists}[B] \rightarrow \text{Lists}[A \times B]$, defined by

$$\text{dist}(a, \langle b_1, b_2, \dots, b_n \rangle) = \langle (a, b_1), (a, b_2), \dots, (a, b_n) \rangle.$$

Basis:

Recursion:

- (c) Give a recursive definition of a function $\text{ins} : \mathbb{R} \times \text{Lists}[\mathbb{R}] \rightarrow \text{Lists}[\mathbb{R}]$ that is supposed to operate as follows: Upon taking a real number x and a list L of real numbers that is ordered in decreasing order, it is supposed to insert x in the list and output a new ordered list with the numbers still in decreasing order. To resolve conflicts the function inserts duplicates on the left of already existing ones.

Basis:

Recursion:

- (d) Give a recursive definition of the function $\text{Apply} : (\mathbb{N} \rightarrow \mathbb{N}) \times \text{Lists}[\mathbb{N}] \rightarrow \text{Lists}[\mathbb{N}]$, defined as follows:

$$\text{Apply}(f, \langle x_0, x_1, \dots, x_n \rangle) = \langle f(x_0), f(x_1), \dots, f(x_n) \rangle.$$

Basis:

Recursion:

- (e) In this part, you may use, if you decide it is convenient to do so, the function Apply that you defined in Part (d), in the spirit of incrementally building computer code by reusing previously defined functions and procedures.

Give a recursive definition of the function

$\text{SpecApply} : (\mathbb{N} \rightarrow \mathbb{N}) \times \text{Lists}[\mathbb{N}] \rightarrow \text{Lists}[\mathbb{N}]$, defined as follows:

$$\text{SpecApply}(f, \langle x_0, x_1, \dots, x_n \rangle) = \langle x_0, f(x_1), f(f(x_2)), f(f(f(x_3))), \dots, f^n(x_n) \rangle.$$

Note that $f^i(x)$ denotes i -fold composition of f with itself and not i -th power.

Basis:

Recursion:

4. (a) A **grammar** is a tuple $G = \langle N, T, S, P \rangle$, where:

(i) N is

(ii) T is

(iii) S is

(iv) P is

(b) Let $G = \langle N, T, S, P \rangle$ be the grammar given in our adopted shorthand notation as follows

$$\begin{aligned} R &\rightarrow AB \\ A &\rightarrow Aa|a \\ B &\rightarrow Bb|\Lambda \end{aligned}$$

Give formally each of the four components of the grammar:

(i) $N =$

(ii) $T =$

(iii) $S =$

(iv) $P =$

(c) The statement “The grammar $S \rightarrow a|SbS$ is ambiguous” is _____

Proof:

(d) Give a grammar for the language $\{a^m b^n : m, n \in \mathbb{N}, n > 0\}$.

(e) Give a grammar for the even palindromes over $\{a, b, c\}$.

(f) Give a grammar for the language $\{a^n b c^n : n \in \mathbb{N}\}^*$.

5. (a) The powerset $\mathcal{P}(\{1, 2\}) = \{ \quad \quad \quad \}$ };
- (b) The powerset $\mathcal{P}(\{0, 1, 2\}) = \{ \quad \quad \quad \}$ };
- (c) Every element in $\mathcal{P}(\{1, 2\})$ “gives rise” to two elements in $\mathcal{P}(\{0, 1, 2\})$. Describe in which way this happens.

- (d) Use the Principle of Induction to prove that the following property $P(n)$ holds for all $n \in \mathbb{N}$:

$$P(n) : \quad 1 + 3 + 3^2 + 3^3 + \cdots + 3^n = \frac{1}{2}(3^{n+1} - 1).$$

Basis:

Induction Hypothesis:

Induction Step:

- (e) (In this part use for your benefit Part (c). Recall $\mathbb{Z}_n = \{0, 1, \dots, n - 1\}$.)
Use the Principle of Induction to prove that the following property $P(n)$ holds for all $n \in \mathbb{N}$:

$$P(n) : \quad |\mathcal{P}(\mathbb{Z}_n)| = 2^n.$$

Basis:

Induction Hypothesis:

Induction Step: